

Original Research

The effects of acute beetroot juice supplementation on lower and upper body isokinetic strength of the wrestlers

Ali TATLICI^{1,*}¹Department of Physical Education and Sport, Faculty of Sport Sciences, Selcuk University, 42130 Konya, Turkey***Correspondence:** alitatlici@selcuk.edu.tr (Ali TATLICI)**Abstract**

Background and objective: Peak torque, as measured by the gold standard method of isokinetic dynamometry, is affected by consumed of dietary nitrate (NO³⁻) in the form of beetroot juice (BRJ). The increasing strength performance effect of BRJ has also started to arouse interest in combat sports. This study aims to investigate the effects of acute dietary nitrate supplementation on lower and upper body isokinetic strength of trained Greco-Roman wrestlers.

Materials and methods: The study is a double-blind, randomized, crossover design in which eight trained male Greco-Roman wrestlers (21.87 ± 2.3 years; height 176.87 cm ± 4.1 cm; body weight 76.75 kg ± 5.4 kg) consumed beetroot juice (140 mL) or placebo (140 mL, cherry juice with lemon juice) 150 min before performing isokinetic strength. Participants performed knee extensors and flexors and shoulder internal and external rotator isokinetic strength test (in the dominant limbs) concentrically at 60° with by Cybex device (Cybex NORM®, Humac, CA, USA). The same trial procedure was repeated with the other supplement (BRJ or Placebo) in a second testing session five days later. BRJ and Placebo trials' isokinetic strength values were compared with parametric testing (paired sample *T*-test).

Results: Although there was a significant difference in peak torque of shoulder internal and external rotator values in favor of BRJ ($P < 0.05$), there was no significant difference in the knee strength test. Additionally, all average isokinetic strength values were significantly increased in the BRJ trial ($P < 0.05$).

Conclusions: These results suggest that BRJ can be used as an ergogenic effect for improving strength values in Greco-Roman wrestling.

Keywords

Beetroot juice; Dietary nitrate; Isokinetic strength; Wrestlers

1. Introduction

Greco-Roman wrestling style is to aim athletes gain superiority over the opponent by holding only the upper body of the opponent with their upper body, whereas freestyle wrestler can use their whole body in the competition [1]. The wrestling rules emphasize aggressive wrestling and scoring by exhibiting repetitive bouts of high-intensity moves (e.g., attacks and counterattacks) instead of blocking or holding the opponent [2, 3]. In wrestling competition, such characteristics as aerobic capacity [4], anaerobic power and maximal

strength are required for success [5]. Wrestlers have to produce high-level maximal strength for defense or offense [6–8]. In wrestling competitions, the upper and lower body maximal strength is one of the most crucial factors to obtain scores by applying the needed technique [9–11]. Studies have emphasized that successful wrestlers are capable of produce higher muscle strength than their less successful counterparts [2, 10–14]. Therefore, increasing muscle strength can help be one of the key factors for wrestlers [1].

Acute dietary supplementation (with beetroot juice) supplementation has been primarily examined in endurance-

based exercise and conflicting results have been found. Compared with endurance-based exercise, beetroot juice (BRJ) effects' on strength were less studied [15]. Studies have shown that BRJ supplementation can affect different muscle fiber types in different level [16]. Since the percentages of skeletal muscle fiber distribution in the body are different [17], it is hypothesized that the effect of BRJ supplementation on lower and upper body strengths in the same individual may be different. To author's knowledge, there are limited studies in the literature investigating acute dietary supplementation to increase wrestlers' muscular strength. BRJ has recently been examined in the literature on different athletes for its potential effects on aerobic capacity, anaerobic power and muscle strength [18].

Beetroot juice, which has a high amount of inorganic nitrate (NO_3^-), has complete attention in sports fields [19]. This NO_3^- in the BRJ is transformed to nitrite (NO_2^-) and then to the nitric oxide (NO) through the loss of oxygen [20]. NO is affecting some physiological processes related to exercise (i.e., regulating blood flow and producing muscle strength) [21]. NO can stimulate the conversion of cyclic guanosine monophosphate (cGMP) that resulting in dilating blood vessels [21], so the amount of blood flow to the muscle tissues increases [22]. Furthermore, the increase of cGMP increases the rate of contraction of muscle fibers and thus may affect maximal strength [21].

The increased NO in blood after drinking BRJ may support the improved type 2 skeletal muscle fiber contractile ability [16] and improved calcium (Ca^{2+}) handling [23], decrease ATP cost of force production [23]. Bender *et al.* [24] found an increased maximal strength in the mid-tight pull. Coggan *et al.* [25] found increased strength in isokinetic dynamometry. Rodríguez-Fernández *et al.* [26] reported that BRJ increased eccentric contractions besides concentric contractions.

The isokinetic dynamometer has high reliability and can measure peak torque at different velocities and throughout the complete range of motion [27–29]. Given that isokinetic dynamometer has been shown as a gold standard strength assessment method [30, 31], isokinetic dynamometry was used in this study. There are studies on the effects of BRJ on the isokinetic strength of the knee [23, 25, 32–40]. The BRJ supplementation effect on isokinetic strength is not well established for the upper and lower body together. However, there are no studies on Greco-Roman wrestlers' upper and lower body isokinetic strength does not exist. Thus, the present study aims to examine the effects of acute BRJ supplementation on upper and lower isokinetic strength parameters in trained Greco-Roman wrestlers.

2. Materials and methods

2.1 Participants

Eight Greco-Roman wrestlers from Konya Selcuklu Municipality wrestling club (age 21.87 ± 2.3 years; height $176.87 \text{ cm} \pm 4.1 \text{ cm}$; body weight $76.75 \text{ kg} \pm 5.4 \text{ kg}$), regularly trained 3 days, volunteered for this study. All participants met the inclusion criteria: being a trained Greco-Roman

wrestler, not having injury or sickness, not taking any other nutritional supplements, and not smoking tobacco in the past six months (Table 1).

Participants were informed about the risks of the study. All participants provided written informed consent. The study was approved by the local ethics committee (Protocol number 79, 19.10.2020, Ethics Committee of Selcuk University, Faculty of Sports Science, Konya, Turkey).

2.2 Experimental design

Participants were asked not to participate in any exercise 24 hours before the testing session and were randomly administered BRJ or placebo (PLA) supplements (four participants in each group). The investigators were not aware of the supplement provided to the participants once provided as the study design was double-blind. Participants waited for 150 min (for peak ergogenic effect) in a lying-down position [41].

Five min standardized warm-up protocol was performed on a cycle ergometer at a rate of 60–70 rpm before isokinetic strength testing. The second trial was conducted with the same frame, with the participants consuming the opposite supplement from the first session. Between the first and second trials, five days waited for washout [42]. Each participant randomly consumed beetroot juice from 2 mL \times 70 mL shots (Beet it, James White Drinks Ltd., Ipswich, UK) or placebo (placebo was made of cherry juice (Cappy, Coca-Cola Company, Istanbul, Turkey)) [43].

Participants consumed a standardized diet (55% carbohydrates, 25% lipids, and 20% protein) 24 hours before both trials. Participants avoid brushing their teeth, using mouthwash or chew gum because of interfering with the effects of BRJ [43]. At last, participants did not eat or drink any foods or drinks that include alcohol, caffeine, or NO_3^- because of the stimulant effects 24 hours prior to testing [44].

2.2.1 Anthropometric measurements

The weight and height of each participant was measured. Body weight was recorded to the nearest 100 g using with scale (Tanita 401 A, Tokyo, Japan) each participant wearing light clothing and no footwear. Height was measured with stadiometer (Holtain Ltd., Crosswell, UK) the nearest $\pm 1 \text{ mm}$ [45].

2.2.2 Isokinetic strength measurement

The isokinetic strength measurements of the knee were performed by an isokinetic dynamometer (Cybex NORM®, Humac, CA, USA) in the kinatropometry laboratory of Selcuk University. Participants were seated in the correct position in the test seat. The participants' holders and the middle sections of the thighs were stabilized to the seat by the tapes. In addition, they were allowed to brace for support by holding the handles on the right and left sides of the seat during the test. The participants were instructed to complete a ROM from 90° to 10° . The point of the beginning was 90° of flexion, then moving into extension.

Each participant was given a familiarization at $60^\circ \text{ sec}^{-1}$ for five repetitions [46]. When the familiarization was done,

TABLE 1. Participants Characteristics.

	N	Min	Max	Mean	Std. dev.
Age	8	20.00	25.00	21.87	2.3
Height	8	171.00	181.00	176.87	4.1
Body Weight	8	71.00	84.00	76.75	5.4

TABLE 2. Comparison of BRJ and PLA Supplementations' Peak Torque.

Parameters	Supplementations	Mean	Std. Deviation	Confidence Interval		P
		(Newton meter)		Lower	Upper	
Extension peak	BRJ	244.25	44.88	-6.54	38.29	0.138
	PLA	228.37	46.46			
Flexion peak	BRJ	141.12	31.25	-12.30	30.55	0.347
	PLA	132.00	25.13			
Internal peak	BRJ	67.37	13.75	0.08	17.41	0.048*
	PLA	58.62	6.41			
External peak	BRJ	47.50	5.63	0.95	9.79	0.024*
	PLA	42.12	4.70			

BRJ, Beetroot juice; PLA, Placebo. * Significant differences ($P < 0.05$).

each participant had a 2 min rest. After the rest period, each participant was asked to perform five repetitions as hard and fast as possible he could at a speed of $60^\circ \text{ sec}^{-1}$. The test for the dominant leg was performed, then each participant was given a 5 min rest, and the shoulder internal and external test started.

Dominant shoulder internal and external rotation strength was obtained from the participants in a standing position, with the elbow flexed at 90° . To measure the muscle strength of the shoulder internal and external, the peak torque (Nm) done with five repetitions at a velocity of $60^\circ \text{ sec}^{-1}$ was determined.

2.3 Statistical analysis

The Shapiro-Wilk test is used to check a data set for normality to make parametric tests applicable. Performance data from the isokinetic strength measurement were analyzed with the paired sample *T*-test (BRJ vs. PLA). Cohen's *d* effect sizes ($d \leq 0.2$, small; $0.5-0.79$, moderate; ≥ 0.8 , large) [47] were calculated to assess the magnitude of difference between experimental trials. All statistical tests were performed using the software package SPSS version 24.0 (SPSS Inc., Chicago, IL, USA). An alpha value of <0.05 was considered to be statistically significant.

3. Results

Peak torque data are shown in Table 2. As intended, BRJ supplementation improved muscle contractile function. Peak torque of knee extension and flexion improved with BRJ ingestion but not statistically. Shoulder internal ($P = 0.048$) (ES = 0.84) and external peak torque ($P = 0.024$) (ES = 1.01) increased significantly in favor of BRJ (large effect size).

Average torque data are shown in Table 3. Average knee strength significantly increased in BRJ trial (extension $P = 0.023$ (ES = 1.02) and flexion $P = 0.027$ (ES = 0.98)). In addition, shoulder internal ($P = 0.023$) (ES = 1.02) and external

average torque ($P = 0.021$) (ES = 1.04) increased significantly in favor of BRJ.

4. Discussion

This study examined the effects of BRJ supplementation on isokinetic strength in trained wrestlers of the upper and lower body. The isokinetic knee extension ($P = 0.13$) and flexion ($P = 0.34$) strength of the wrestlers did not change significantly in both trials, but average strength increased in the BRJ trial than the PLA trial (extension 9.57% and flexion 12.94%). Furthermore, the peak isokinetic strength of the shoulder increased in favor of the BRJ trial (internal 15.06% and external torque 13.39%). The average isokinetic shoulder internal (19.61%) and external (12.81%) strength of the wrestlers improved significantly.

The supplementation of BRJ increases NO availability in metabolism [37]. Literature shows that NO affects contractile properties in animals, also in human muscle [23-25, 33, 37, 48]. However, the present study found no significant increased knee isokinetic strength in both extension and flexion parameters at $60^\circ \text{ sec}^{-1}$ (slow angular velocity), consistent with Jonvik *et al.* [21] and Kokkinoplitis *et al.*'s findings [39]. In addition, studies showed that peak isokinetic strength of the knee values performed at $90^\circ \text{ sec}^{-1}$ angular velocity (moderate angular velocity) was not significantly affected by NO^{-3} supplementation [25, 37, 40]. Inconsistent with the present study, Jonvik *et al.* [21] found significant improvement in knee flexion at $60^\circ \text{ sec}^{-1}$ angular velocity, but their participants were formed by recreationally active individuals. Individuals with more training benefit less from NO^{-3} supplementation [41, 49], which could be the underlying reason why isokinetic leg flexor peak strength was not significant in our study. In this study, BRJ supplementation did not significantly affect the peak isokinetic strength of the lower body but increased the peak isokinetic strength of the upper body significantly. NO^{3-} can be more effective at

TABLE 3. Comparison of BRJ and PLA Supplementations' Average Torque.

Parameters	Supplementations	Mean (Watts)	Std. Deviation	Confidence Interval		P
				Lower	Upper	
Extension average	BRJ	170.50	25.57	2.64	25.60	0.023*
	PLA	156.37	25.98			
Flexion average	BRJ	107.75	18.69	1.95	23.79	0.027*
	PLA	94.87	16.03			
Internal average	BRJ	55.12	11.60	1.69	16.55	0.023*
	PLA	46.00	5.23			
External average	BRJ	38.62	5.70	0.85	7.64	0.021*
	PLA	34.37	4.53			

* Significant differences ($P < 0.05$).

type 2 muscle fiber contraction [50]. Horswill *et al.* [51] reported that Greco-Roman wrestlers could generate more anaerobic power by the upper body than lower due to type II muscle fiber distribution. It can be hypothesized that type 2 muscle fiber distribution can be dominant in the shoulder muscle group than leg muscle, therefore, shoulder internal and external peak torque increased significantly in this study. To the author's knowledge, although there is no study on upper extremity and isokinetic BRJ-strength in the literature, Williams *et al.* [15] reported that bench press power positively increased. Similar to our findings obtained in this study, there are studies in the literature that emphasized that lower extremity average strength outputs improved after ingestion of BRJ [26, 35, 52]. BRJ supplementation reported declining the ATP cost of force production [21, 52]. Since local perfusion, fatigue resistance, increasing the intracellular release of calcium in the muscle and contractility of fast-twitch muscle fibers, in particular, can be improved by BRJ supplementation [53], the isokinetic strengths outputs were thought to increase in this study.

Due to the nature of wrestling, wrestlers should have advanced maximal strength, strength endurance, aerobic capacity and anaerobic power levels in addition to technical and tactical skills [1, 5]. To perform a proper technique during the wrestling match, it is often necessary to move with high maximal strength and be able to repeat that movement at the highest strength possible [1]. Furthermore, the wrestlers should maintain maximal strength when they eat less for rapidly losing weight because studies stated that the muscle strength [54] and power [55] of the wrestlers could be affected by the rapid weight loss [56]. Thus, increasing maximal strength and strength endurance with acute BRJ supplementation can help wrestler's performance in competition and training.

This study was conducted with limited resources, and it only provided insights into the strength of the lower and upper body of Greco-Roman wrestlers. Another limitation of the present study can be said as the number of participants was eight.

5. Conclusions

The data obtained by this study indicates that BRJ supplementation increased peak and average isokinetic

strength of the lower and upper body of trained Greco-Roman wrestlers. These results suggest that beetroot juice can be used as an ergogenic aid before competition for increasing muscle strength performance in trained Greco-Roman wrestlers.

Ethics approval and consent to participate

The study was approved by the local ethics committee (Protocol number 79, 19.10. 2020, Ethics Committee of Selcuk University, Faculty of Sports Science, Konya, Turkey). Before the assessment, every participant received the same detailed information about the testing procedure. Every participant signed the informed consent.

Acknowledgment

Thanks to all the peer reviewers for their opinions and suggestions.

Funding

This research received no external funding.

Conflict of interest

The author declares no conflict of interest.

References

- [1] Chaabene H, Negra Y, Bouguezzi R, Mkaouer B, Franchini E, Julio U, *et al.* Physical and physiological attributes of wrestlers: an update. *Journal of Strength and Conditioning Research.* 2017; 31: 1411–1442.
- [2] Yoon J. Physiological profiles of elite senior wrestlers. *Sports Medicine.* 2002; 32: 225–233.
- [3] Horswill CA. Applied physiology of amateur wrestling. *Sports Medicine.* 1992; 14: 114–143.
- [4] Alexandrova A, Eroglu Y, Petrov L, Makaveev R, Georgieva A, Tzvetanova E. Blood plasma oxidative stress parameters after maximal oxygen consumption test in wrestlers. *International Journal of Sport Studies.* 2016; 6: 359–366.
- [5] Demirkan E, Koz M, Kutlu M, Favre M. Comparison of physical and physiological profiles in elite and amateur young wrestlers. *Journal of Strength and Conditioning Research.* 2015; 29: 1876–1883.
- [6] Mirzaei B, Curby DG, Barbas I, Lotfi N. Physical fitness measures of cadet wrestlers. *International Journal of Wrestling Science.* 2011; 1: 63–66.

- [7] Passelergue PA, Lac G. Salivary hormonal responses and performance changes during 15 weeks of mixed aerobic and weight training in elite junior wrestlers. *Journal of Strength and Conditioning Research*. 2012; 26: 3049–3058.
- [8] Mirzaei B, Curby DG, Rahmani-Nia F, Moghadasi M. Physiological profile of elite Iranian junior freestyle wrestlers. *Journal of Strength and Conditioning Research*. 2009; 23: 2339–2344.
- [9] Mirzaei B, Curby DG, Barbas I, Lotfi N. Anthropometric and physical fitness traits of four-time World Greco-Roman wrestling champion in relation to national norms: a case study. *Journal of Human Sport and Exercise*. 2011; 6: 406–413.
- [10] Pallarés JG, López-Gullón JM, Torres-Bonete MD, Izquierdo M. Physical fitness factors to predict female Olympic wrestling performance and sex differences. *Journal of Strength and Conditioning Research*. 2012; 26: 794–803.
- [11] Morán-Navarro R, Valverde-Conesa A, López-Gullón JM, la Cruz-Sánchez D, Pallarés JG. Can balance skills predict Olympic wrestling performance? *Journal of Sport & Health Research*. 2015; 7: 19–30.
- [12] He ZH, Feng LS, Zhang HJ, Xu KY, Chi FT, Tao DL, *et al.* Physiological profile of elite Chinese female wrestlers. *Journal of Strength and Conditioning Research*. 2013; 27: 2374–2395.
- [13] Zhang CG, Xiu ZT, He ZH, Lu DL, Tao DL, Tang TM, *et al.* Investigation on sports injuries in Chinese elite wrestlers. *J Tianjin Inst Phys Educ*. 2005; 3: 69–71.
- [14] Horswill CA, Scott, JR, Galea P. Comparison of maximum aerobic power, maximum anaerobic power, and skinfold thickness of elite and nonelite junior wrestlers. *International Journal of Sports Medicine*. 1989; 10: 165–168.
- [15] Williams TD, Martin MP, Mintz JA, Rogers RR, Ballmann CG. Effect of acute beetroot juice supplementation on bench press power, velocity, and repetition volume. *Journal of Strength and Conditioning Research*. 2020; 34: 924–928.
- [16] Jones AM. Dietary nitrate supplementation and exercise performance. *Sports Medicine*. 2014; 44: 35–45.
- [17] Gollnick PD, Armstrong RB, Saubert CW, Piehl K, Saltin B. Enzyme activity and fiber composition in skeletal muscle of untrained and trained men. *Journal of Applied Physiology*. 1972; 33: 312–319.
- [18] Zamani H, de Joode MEJR, Hosseini IJ, Henckens NFT, Guggeis MA, Berends JE, *et al.* The benefits and risks of beetroot juice consumption: a systematic review. *Critical Reviews in Food Science and Nutrition*. 2021; 61: 788–804.
- [19] Larsen FJ, Weitzberg E, Lundberg JO, Ekblom B. Dietary nitrate reduces maximal oxygen consumption while maintaining work performance in maximal exercise. *Free Radical Biology and Medicine*. 2010; 48: 342–347.
- [20] Domínguez R, Cuenca E, Maté-Muñoz J, García-Fernández P, Serrapaya N, Estevan M, *et al.* Effects of beetroot juice supplementation on cardiorespiratory endurance in athletes. A Systematic Review. *Nutrients*. 2017; 9: 43.
- [21] Jonvik KL, Hoogervorst D, Peelen HB, de Niet M, Verdijk LB, van Loon LJC. The impact of beetroot juice supplementation on muscular endurance, maximal strength and countermovement jump performance. *European Journal of Sport Science*. 2021; 21: 871–878.
- [22] Bonilla Ocampo DA, Paipilla AF, Marín E, Vargas-Molina S, Petro JL, Pérez-Idárraga A. Dietary nitrate from beetroot juice for hypertension: a systematic review. *Biomolecules*. 2018; 8: 134.
- [23] Whitfield J, Gamu D, Heigenhauser GJ, Van Loon LJC, Spriet LL, Tupling AR, *et al.* Beetroot juice increases human muscle force without changing Ca²⁺-handling proteins. *Medicine & Science in Sports & Exercise*. 2017; 49: 2016–2024.
- [24] Bender D, Townsend JR, Vantrease WC, Marshall AC, Henry RN, Heffington SH, *et al.* Acute beetroot juice administration improves peak isometric force production in adolescent males. *Applied Physiology, Nutrition, and Metabolism*. 2018; 43: 816–821.
- [25] Coggan AR, Leibowitz JL, Spearie CA, Kadkhodayan A, Thomas DP, Ramamurthy S, *et al.* Acute dietary nitrate intake improves muscle contractile function in patients with heart failure. *Circulation: Heart Failure*. 2015; 8: 914–920.
- [26] Rodríguez-Fernández A, Castillo D, Raya-González J, Domínguez R, Bailey SJ. Beetroot juice supplementation increases concentric and eccentric muscle power output. Original investigation. *Journal of Science and Medicine in Sport*. 2021; 24: 80–84.
- [27] Lago-Rodríguez Á, Domínguez R, Ramos-Álvarez JJ, Tobal FM, Jodra P, Tan R, *et al.* The effect of dietary nitrate supplementation on isokinetic torque in adults: a systematic review and meta-analysis. *Nutrients*. 2020; 12: 3022.
- [28] Mueller S, Stoll J, Mueller J, Mayer F. Validity of isokinetic trunk measurements with respect to healthy adults, athletes and low back pain patients. *Isokinetics and Exercise Science*. 2012; 20: 255–266.
- [29] Ozdemir C, Oztügen K, Sarpel T, Günası O, Bildircin C, Coşkun F, Kurdak SS. Monitoring the improvement of muscle performance using isokinetic dynamometry: A comparative analysis based on the full range of motion vs. the valid isokinetic sector. *Isokinetics and Exercise Science*. 2018; 26: 257–263.
- [30] Kristensen O H, Stenager E, Dalgas U. Muscle strength and poststroke hemiplegia: a systematic review of muscle strength assessment and muscle strength impairment. *Archives of Physical Medicine and Rehabilitation*. 2017; 98: 368–380.
- [31] Muñoz-Bermejo L, Pérez-Gómez J, Manzano F, Collado-Mateo D, Villafaina S, Adsuar JC. Reliability of isokinetic knee strength measurements in children: A systematic review and meta-analysis. *PLoS ONE*. 2019; 14: e0226274.
- [32] Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, *et al.* Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *Journal of Applied Physiology*. 2010; 109: 135–148.
- [33] HAIDER G, FOLLAND JP. Nitrate supplementation enhances the contractile properties of human skeletal muscle. *Medicine & Science in Sports & Exercise*. 2014; 46: 2234–2243.
- [34] Hoon MW, Fornusek C, Chapman PG, Johnson NA. The effect of nitrate supplementation on muscle contraction in healthy adults. *European Journal of Sport Science*. 2015; 15: 712–719.
- [35] Tillin NA, Moudy S, Nourse KM, Tyler CJ. Nitrate supplement benefits contractile forces in fatigued but not unfatigued muscle. *Medicine & Science in Sports & Exercise*. 2018; 50: 2122–2131.
- [36] Fulford J, Winyard PG, Vanhatalo A, Bailey SJ, Blackwell JR, Jones AM. Influence of dietary nitrate supplementation on human skeletal muscle metabolism and force production during maximum voluntary contractions. *Pflügers Archiv - European Journal of Physiology*. 2013; 465: 517–528.
- [37] Coggan AR, Leibowitz JL, Kadkhodayan A, Thomas DP, Ramamurthy S, Spearie CA, *et al.* Effect of acute dietary nitrate intake on maximal knee extensor speed and power in healthy men and women. *Nitric Oxide*. 2015; 48: 16–21.
- [38] Coggan AR, Hoffman RL, Gray DA, Moorthi RN, Thomas DP, Leibowitz JL, *et al.* A single dose of dietary nitrate increases maximal knee extensor angular velocity and power in healthy older men and women. *The Journals of Gerontology: Series A*. 2020; 75: 1154–1160.
- [39] Kokkinoplitis K, Chester N. The effect of beetroot juice on repeated sprint performance and muscle force production. *Journal of Physical Education and Sport*. 2014; 14: 242–247.
- [40] Coggan AR, Broadstreet SR, Mikhalkova D, Bole I, Leibowitz JL, Kadkhodayan A, *et al.* Dietary nitrate-induced increases in human muscle power: high versus low responders. *Physiological Reports*. 2018; 6: e13575.
- [41] Wilkerson DP, Hayward GM, Bailey SJ, Vanhatalo A, Blackwell JR, Jones AM. Influence of acute dietary nitrate supplementation on 50 mile time trial performance in well-trained cyclists. *European Journal of Applied Physiology*. 2012; 112: 4127–4134.
- [42] Rimer EG, Peterson LR, Coggan AR, Martin JC. Acute dietary nitrate supplementation increases maximal cycling power in athletes. *International Journal of Sports Physiology and Performance*. 2016; 11: 715–720.
- [43] Tatlici, A, Çakmakçi, O. The effects of acute dietary nitrate supplementation on anaerobic power of elite boxers. *Medicina Dello Sport*. 2019; 72: 225–233.
- [44] Lansley KE, Winyard PG, Fulford J, Vanhatalo A, Bailey SJ, Blackwell

- JR, *et al.* Dietary nitrate supplementation reduces the O₂ cost of walking and running: a placebo-controlled study. *Journal of Applied Physiology*. 2011; 110: 591–600.
- [45] Zaekov N, Petrov L, Rankov K, Nikolova M, Makaveev R, Nikolov N. Determining energy expenditure from heart rate monitoring during training sessions at bulgarian wrestlers. *International Journal Bioautomation*. 2019; 23: 251–258.
- [46] Van Cingel EHR, Kleinrensink GJ, Rooijens PPGM, Uitterlinden EJ, Aufdemkampe G, Stoeckart R. Learning effect in isokinetic testing of ankle invertors and evertors. *Isokinetics and Exercise Science*. 2001; 9: 171–177.
- [47] Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Academic Press: New York. 2013.
- [48] Wickham KA, McCarthy DG, Pereira JM, Cervone DT, Verdijk LB, van Loon LJC, *et al.* No effect of beetroot juice supplementation on exercise economy and performance in recreationally active females despite increased torque production. *Physiological Reports*. 2019; 7: e13982.
- [49] Christensen PM, Nyberg M, Bangsbo J. Influence of nitrate supplementation on VO₂ kinetics and endurance of elite cyclists. *Scandinavian Journal of Medicine & Science in Sports*. 2013; 23: e21–e31.
- [50] Hernández A, Schiffer TA, Ivarsson N, Cheng AJ, Bruton JD, Lundberg JO, *et al.* Dietary nitrate increases tetanic [Ca²⁺]_i and contractile force in mouse fast-twitch muscle. *The Journal of Physiology*. 2012; 590: 3575–3583.
- [51] Horswill CA, Miller JE, Scott JR, Smith CM, Welk G, Van Handel P. Anaerobic and aerobic power in arms and legs of elite senior wrestlers. *International Journal of Sports Medicine*. 1992; 13: 558–561.
- [52] Jones AM, Ferguson SK, Bailey SJ, Vanhatalo A, Poole DC. Fiber type-specific effects of dietary nitrate. *Exercise and Sport Sciences Reviews*. 2016; 44: 53–60.
- [53] Houston ME, Marrin DA, Green HJ, Thomson JA. The effect of rapid weight loss on physiological functions in wrestlers. *the Physician and Sportsmedicine*. 1981; 9: 73–78.
- [54] Kraemer WJ, Fry AC, Rubin MR, Triplett-Mcbride T, Gordon SE, Perry Koziris L, *et al.* Physiological and performance responses to tournament wrestling. *Medicine and Science in Sports and Exercise*. 2001; 33: 1367–1378.
- [55] Rankin JW, Ocel JV, Craft LL. Effect of weight loss and refeeding diet composition on anaerobic performance in wrestlers. *Medicine and Science in Sports and Exercise*. 1996; 28: 1292–1299.
- [56] Marttinen RHJ, Judelson DA, Wiersma LD, Coburn JW. Effects of self-selected mass loss on performance and mood in collegiate wrestlers. *Journal of Strength and Conditioning Research*. 2011; 25: 1010–1015.